ELSEVIER

Journal of Affective Disorders 80 (2004) 303–305

JOURNAL OF AFFECTIVE DISORDERS

www.elsevier.com/locate/jad

Contents (Vol. 80)

Issue 1 May 2004

Research reports

Bipolar disorder among an isolated island community in Ethiopia
A. Fekadu, T. Shibre, A. Alem, D. Kebede, S. Kebreab, A. Negash and M.J. Owen (UK, Ethiopia) 1

Differential response of bipolar and normal control lymphoblastoid cell sodium pump to ethacrynic acid
R. Li and R.S. El-Mallakh (USA) 11

Psychotic symptoms in pediatric bipolar disorder
M.N. Pavuluri, E.S. Herbener and J.A. Sweeney (USA) 19

A comparison of the phenomenology and treatment of youths and adults with bipolar I disorder in a state mental health system
J.M. Jerrell and M.A. Shugart (USA) 29

Identifying depression in the first postpartum year: guidelines for office-based screening and referral
K.S. Peindl, K.L. Wisner and B.H. Hanusa (USA) 37

Single photon emission computed tomography in posttraumatic stress disorder before and after treatment with a selective serotonin reuptake inhibitor
S. Seedat, J. Warwick, B. van Heerden, C. Hugo, N. Zungu-Dirwayi, J. Van Kradenburg and D.J. Stein (South Africa) 45

Correlation between cerebral blood flow and items of the Hamilton Rating Scale for Depression in antidepressant-naive patients
A. Graff-Guerrero, J. González-Olvera, Y. Mendoza-Espinosa, V. Vaugier and J.C. García-Reyna (Mexico) 55

The course of anxiety and depression through pregnancy and the postpartum in a community sample
J. Heron, T.G. O'Connor, J. Evans, J. Golding, V. Glover and the ALSPAC Study Team (UK, USA) 65

Brief reports

Personality disorder and social anxiety predict delayed response in drug and behavioral treatment of panic disorder
P. Berger, G. Sachs, M. Amering, A. Holzinger, B. Bankier and H. Katschnig (Austria) 75

An illustration of how a self-report diagnostic screening scale could improve the internal validity of antidepressant efficacy trials
M. Zimmerman, I. Chelminski and M. Posternak (USA) 79

Preliminary communication

Is there evidence for a male depressive syndrome in inpatients with major depression?
A. Maria Möller-Leimkühler, R. Bottlender, A. Strauß and W. Rutz (Germany, Denmark) 87

doi:10.1016/S0165-0327(04)00136-3

Letter to the Editor

Letter to the Editor
S.M. Hunt (UK) 95

Instructions to Authors 97

Issues 2–3 June 2004

Review

Psychotherapy of bipolar disorder: a review
S. Jones (UK) 101

Special article

Capacity to consent to research among patients with bipolar disorder
S. Misra and L. Ganzini (USA) 115

Research reports

Heritability of individual depressive symptoms
K.L. Jang, W.J. Livesley, S. Taylor, M.B. Stein and E.C. Moon (Canada, USA) 125
Residual symptoms at remission from depression: impact on long-term outcome
N. Kennedy and E.S. Paykel (UK) 135
The sleep of remitted bipolar outpatients: a controlled naturalistic study using actigraphy
A. Millar, C.A. Espie and J. Scott (UK) 145
Axis I and II comorbidity in a large sample of patients with obsessive–compulsive disorder
D. Denys, N. Tenney, H.J.G.M. van Megen, F. de Geus and H.G.M. Westenberg (The Netherlands) 155
Comorbidity of generalized social anxiety disorder and depression in a pediatric primary care sample
D.A. Chavira, M.B. Stein, K. Bailey and M.T. Stein (USA) 163
Are effects of depression management training for General Practitioners on patient outcomes mediated by improvements in the process of care?
T.W.D.P. van Os, R.H.S. van den Brink, B.G. Tiemens, J.A. Jenner, K. van der Meer and J. Ormel (The Netherlands) 173
An examination of seasonality experienced by Australians living in a continental temperate climate zone
R.A. Parslow, A.F. Jorm, P. Butterworth, P.A. Jacomb and B. Rodgers (Australia) 181
The process of suicidal planning among medical doctors: predictors in a longitudinal Norwegian sample
R. Tyssen, E. Hem, P. Vaglum, N.T. Grønvold and Ø. Ekeberg (Norway) 191
Depression, partnership, social support, and parenting: interaction of maternal factors with behavioral problems of the child
J.E. Herwig, M. Wirtz and J. Bengel (Germany) 199
The reliability and validity of the Seasonal Pattern Assessment Questionnaire: a comparison between patient groups
P.P.A. Mersch, N.C. Vastenburg, Y. Meesters, A.L. Bouhuys, D.G.M. Beersma, R.H. van den Hoofdakker and J.A. den Boer (The Netherlands) 209
Neurological soft signs in bipolar I disorder patients
A. Negash, D. Kebede, A. Alem, Z. Melaku, N. Deyessa, T. Shibire, A. Fekadu, D. Fekadu, L. Jacobsson and G. Kullgren (Ethiopia, Sweden) 221

Comorbidity of PTSD and depression: associations with trauma exposure, symptom severity and functional impairment in Bosnian refugees resettled in Australia
S. Momartin, D. Silove, V. Manicavasagar and Z. Steel (Australia) 231
What depressive symptoms are associated with the use of care services? Results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS)
M. ten Have, R. de Graaf, W. Vollebergh and A. Beekman (The Netherlands) 239
Increased (*E*)-4-hydroxy-2-nonenal and asymmetric dimethylarginine concentrations and decreased nitric oxide concentrations in the plasma of patients with major depression
M.L. Selley (Australia) 249

Brief reports
Worsening of depressive symptoms 6 months after an acute coronary event in older adults is associated with impairment of cardiac autonomic function
M.S.L. de Guevara, S.I. Schauffele, L.C. Nicola-Siri, R.D. Fahrer, E. Ortíz-Frágola, J.A. Martínez-Martínez, D.P. Cardinali and S.M. Guinjoan (Argentina) 257
The use of the GBI in a population of adolescent offspring of parents with a bipolar disorder
C.G. Reichart, J. van der Ende, M. Wals, M.H.J. Hillegers, J. Ormel, W.A. Nolen and F.C. Verhulst (The Netherlands) 263
Controlled lithium discontinuation in bipolar patients with good response to long-term lithium prophylaxis
O. Yazıcı, K. Kora, A. Polat and M. Şaylan (Turkey) 269

Preliminary communications
Cognitive-behavioral therapy, light therapy, and their combination in treating seasonal affective disorder
K.J. Rohan, K.T. Lindsey, K.A. Roecklein and T.J. Lacy (USA) 273
Sensitivity to cholecystokinin-tetrapeptide in major depression
D. Koszycki, J. Copen and J. Bradwejn (Canada) 285

Letter to the Editor
Biological markers and psychophysiological methods as long term response predictors to fluoxetine treatment
K.N. Fountoulakis, S. Kaprinis, A. Iacovides, F. Fotiou and G. Kaprinis (Greece) 291

Instructions to Authors 293

Author index 295

Subject index 299

Contents (Vol. 80) 303

Are effects of depression management training for General Practitioners on patient outcomes mediated by improvements in the process of care?
T.W.D.P. van Os, R.H.S. van den Brink, B.G. Tiemens, J.A. Jenner, K. van der Meer and J. Ormel (The Netherlands) 173

An examination of seasonality experienced by Australians living in a continental temperate climate zone
R.A. Parslow, A.F. Jorm, P. Butterworth, P.A. Jacomb and B. Rodgers (Australia) 181

The process of suicidal planning among medical doctors: predictors in a longitudinal Norwegian sample
R. Tyssen, E. Hem, P. Vaglum, N.T. Grønvold and Ø. Ekeberg (Norway) 191

Depression, partnership, social support, and parenting: interaction of maternal factors with behavioral problems of the child
J.E. Herwig, M. Wirtz and J. Bengel (Germany) 199

The reliability and validity of the Seasonal Pattern Assessment Questionnaire: a comparison between patient groups
P.P.A. Mersch, N.C. Vastenburg, Y. Meesters, A.L. Bouhuys, D.G.M. Beersma, R.H. van den Hoofdakker and J.A. den Boer (The Netherlands) 209

Neurological soft signs in bipolar I disorder patients
A. Negash, D. Kebede, A. Alem, Z. Melaku, N. Deyessa, T. Shibire, A. Fekadu, D. Fekadu, L. Jacobsson and G. Kullgren (Ethiopia, Sweden) 221

Comorbidity of PTSD and depression: associations with trauma exposure, symptom severity and functional impairment in Bosnian refugees resettled in Australia
S. Momartin, D. Silove, V. Manicavasagar and Z. Steel (Australia) 231

What depressive symptoms are associated with the use of care services? Results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS)
M. ten Have, R. de Graaf, W. Vollebergh and A. Beekman (The Netherlands) 239

Increased (*E*)-4-hydroxy-2-nonenal and asymmetric dimethylarginine concentrations and decreased nitric oxide concentrations in the plasma of patients with major depression
M.L. Selley (Australia) 249

Brief reports

Worsening of depressive symptoms 6 months after an acute coronary event in older adults is associated with impairment of cardiac autonomic function
M.S.L. de Guevara, S.I. Schauffele, L.C. Nicola-Siri, R.D. Fahrer, E. Ortíz-Frágola, J.A. Martínez-Martínez, D.P. Cardinali and S.M. Guinjoan (Argentina) 257

The use of the GBI in a population of adolescent offspring of parents with a bipolar disorder
C.G. Reichart, J. van der Ende, M. Wals, M.H.J. Hillegers, J. Ormel, W.A. Nolen and F.C. Verhulst (The Netherlands) 263

Controlled lithium discontinuation in bipolar patients with good response to long-term lithium prophylaxis
O. Yazıcı, K. Kora, A. Polat and M. Şaylan (Turkey) 269

Preliminary communications

Cognitive-behavioral therapy, light therapy, and their combination in treating seasonal affective disorder
K.J. Rohan, K.T. Lindsey, K.A. Roecklein and T.J. Lacy (USA) 273

Sensitivity to cholecystokinin-tetrapeptide in major depression
D. Koszycki, J. Copen and J. Bradwejn (Canada) 285

Letter to the Editor

Biological markers and psychophysiological methods as long term response predictors to fluoxetine treatment
K.N. Fountoulakis, S. Kaprinis, A. Iacovides, F. Fotiou and G. Kaprinis (Greece) 291

Instructions to Authors 293

Author index 295

Subject index 299

Contents (Vol. 80) 303